arXiv:q-alg/9708027v1 23 Aug 1997

TOPICS IN ISOTOPIC PAIRS AND THEIR REPRESENTATIONS. III. BUNCHES OF LIE ALGEBRAS AND MODIFIED CLASSICAL YANG-BAXTER EQUATION

D.V.Juriev[1]

Research Center for Mathematical Physics and Informatics “Thalassa Aitheria”,
ul.Miklukho-Maklaya 20-180, Moscow 117437 Russia.
E-mail: denis@juriev.msk.ru

q-alg/9708027

This short note is a brief comment to certain aspects of the theory of classical r-matrix [1] and bihamiltonian formalism [2], which motivations lie in constructions of articles [3].

1. Linear bunches of Lie algebras and Lie $\widetilde{\mathrm{m}}\mathrm{YB}$-algebras

This paragraph is devoted to the anlysis of some algebraic structures associated with linear bunches of Lie algebras.

1.1. Γ-bunches of Lie algebras and tangent bracket.

Definition 1A. The *(one-parametric) bunch of Lie algebras* is a linear space $\mathfrak{g}$ supplied by a family of Lie brackets $[\cdot,\cdot]_\lambda$ ($\lambda\in\mathbb{R}$). The Lie algebra from $\mathfrak{g}$, which is defined by the bracket $[\cdot,\cdot]_\lambda$, is denoted by $\mathfrak{g}_\lambda$. A bunch of Lie algebras $\mathfrak{g}$ is called the Γ*-bunch* if there are defined homomorphisms R_λ from the Lie algebras $\mathfrak{g}_\lambda$ into the Lie algebra $\mathfrak{g}_0$, otherwords, the equality

$$R_\lambda[X,Y]_\lambda=[R_\lambda X,R_\lambda Y],$$

where $[\cdot,\cdot]=[\cdot,\cdot]_0$, holds for all X and Y from $\mathfrak{g}$.

Below there will be considered only smooth bunches, i.e. the bunches, which Lie brackets form a smooth family, and smooth Γ-bunches, for which the family of homomorphisms R_λ is also smooth.

---

[1] This is an English translation of the original Russian version, which is located at the end of the article as an appendix. In the case of any differences between English and Russian versions caused by a translation the least has the priority as the original one.

Typeset by $\mathcal{A}_{\mathcal{M}}\mathcal{S}$-TEX

Proposition 1. *Let $\mathfrak{g}$ be a smooth Γ-bunch of Lie algebras. Define the tangent bracket $[\cdot,\cdot]_R$ as*

$$[X,Y]_R = \left.\frac{d[X,Y]_\lambda}{d\lambda}\right|_{\lambda=0}.$$

Then

$$[X,Y]_R = [RX,Y] + [X,RY] - R[X,Y],$$

where

$$R = \left.\frac{dR_\lambda}{d\lambda}\right|_{\lambda=0}.$$

Remark 1. The bracket $[\cdot,\cdot]_R$ is trivial (is identical zero) if and only if R is a derivative of the bracket $[\cdot,\cdot]$.

Remark 2. The bracket $[\cdot,\cdot]_R$ is a Lie one if and only if

$$\big([B(X,Y),Z] + B([X,Y],Z)\big) + c.p. = 0,$$

where

$$B(X,Y) = R[RX,Y] + R[X,RY] - [RX,RY] - R^2[X,Y].$$

The expanded expression for $B(X,Y)$ may be compactified into the binomial formula

$$B(X,Y) = R[X,Y]_R - [RX,RY].$$

1.2. Linear Γ-bunches of Lie algebras and the modified Yang-Baxter equation. Note that the bracket $[\cdot,\cdot]$ and the tangent bracket $[\cdot,\cdot]_R$ in any Γ-bunch of Lie algebras always obey the identity

$$\big([[X,Y]_R,Z] + [[X,Y],Z]_R\big) + c.p. = 0.$$

Hence, if the tangent bracket $[\cdot,\cdot]_R$ is a Lie one, then it is compatible with the bracket $[\cdot,\cdot]$. This circumstance motivates to consider the linear Γ-bunches of Lie algebras.

A bunch of Lie algebras $\mathfrak{g}$ is called linear if the Lie brackets $[\cdot,\cdot]_\lambda$ from a linear family, otherwords, $[\cdot,\cdot]_\lambda = (1-\lambda)[\cdot,\cdot]+\lambda[\cdot,\cdot]_1$. A Γ-bunch of of Lie algebras is called linear if not only Lie brackets but also the homomorphisms R_λ form a linear family, i.e. $R_\lambda = 1 + \lambda R$. The condition of linearity of the family of homomorphisms R_λ is natural, because for any nonlinear family R_λ its linear part $1+\lambda R$ ($R = \frac{dR_\lambda}{d\lambda}\big|_{\lambda=0}$) defines the same linear family of brackets $[\cdot,\cdot]_\lambda$.

Theorem 1A. *For a linear Γ-bunch of Lie algebras $\mathfrak{g}$ the tangent bracket $[\cdot,\cdot]_R$ (which is a Lie one in this case) obeys the following condition*

$$R[X,Y]_R = [RX,RY],$$

otherwords,

$$R[RX,Y] + R[X,RY] = [RX,RY] + R^2[X,Y].$$

The theorem means that the condition that the bracket $[\cdot,\cdot]_R$ is a Lie one from remark 2 is factorized in case of linear Γ-bunches and is reduced to the claim of the identical equality of the expression $B(X,Y)$ to zero. Note that the fact that the bracket $[\cdot,\cdot]_R$ is a Lie one in view of its compatibility with $[\cdot,\cdot]$ means that the bunch $[\cdot,\cdot]+\lambda[\cdot,\cdot]_R$ is a linear bunch of Lie algebras, however, it is not obligatory a Γ-bunch, therefore the condition of remark 2 is not factorized generally.

The least identity of theorem 1A coincides with the modified classical Yang-Baxter equation [1] if $R^2=1$ that holds for the most important examples considered in [1]. In this case as it was shown in [1] the bracket $[\cdot,\cdot]'_R$ defined as

$$[X,Y]'_R=[RX,Y]+[X,RY]=[X,Y]_R+R[X,Y]$$

is a Lie one. The general condition that the bracket $[\cdot,\cdot]'_R$ is a Lie one has the form

$$[R^2[X,Y],Z]+c.p.=0.$$

The bracket $[\cdot,\cdot]'_R$ plays a crucial role in the factorization method [1]. Let us consider an example when $R^2\neq 1$, nevertheless, $[R^2[X,Y],Z]+c.p.=0$.

Example 1. Let us consider the Witt algebra with the basis e_k ($k\in\mathbb{Z}$) and the commutation relations

$$[e_i,e_j]=(i-j)e_{i+j}.$$

Introduce the operators R_n of the form

$$R_n(e_i)=e_{i+n}.$$

Each operator R_n defines a structure of $\widetilde{\mathrm{m}}$YB-algebra on the Witt algebra, for none n the squares of operators R_n are not equal to the identity but for all n the equality $[R_n^2[X,Y],Z]+c.p.=0$ holds.

Remark 3. The linear Γ-bunches of Lie algebras do not form a submanifold (in sense of [4,5]) of the manifold of all linear bunches of Lie algebras. The identities, which connect the brackets $[\cdot,\cdot]$ and $[\cdot,\cdot]_R$ (as well as $[\cdot,\cdot]$ and $[\cdot,\cdot]'_R$) are not known.

Definition 1B. A linear bunch of Lie algebras $\mathfrak{g}$ is called the *linear* $\Pi\Gamma$*-bunch* if it admits a monomorphism into a linear Γ-bunch of Lie algebras.

As it was marked above a description of the class of all linear $\Pi\Gamma$-bunches of Lie algebras by the complete system of identities is not known.

Let us consider an example of the linear $\Pi\Gamma$-bunch of Lie algebras, whcih is not a linear Γ-bunch.

Example 2 (cf.[3]). Let $\mathfrak{g}=\mathfrak{sa}(n)$ be the Lie algebra of all skew-symmetric $n\times n$ matrices with the standard commutator $[X,Y]=XY-YX$ and Q be an arbitrary nonscalar symmetric $n\times n$ matrix. Define a family of compatible Lie brackets $[\cdot,\cdot]_\lambda$ as

$$[X,Y]_\lambda=[X,Y]+\lambda(XQY-YQX).$$

The obtained linear bunch of Lie algebras is not a linear Γ-bunch, because in view of the nonscalarity of the matrix Q some of the algebras $\mathfrak{g}_\lambda$ being isomorphic to $\mathfrak{sa}(p,q)$ do not admit any homomorphisms into $\mathfrak{sa}(n)$. On the other hand this bunch may be imbed into the bunch of all $n\times n$ matrices with the same brackets, which is a linear Γ-bunch with homomorphisms R_λ defined by the operators of the left or right multiplication on the matrices $Q_\lambda=1+\lambda Q$.

1.3. Lie $\widetilde{\mathrm{m}}$YB-algebras and linear Γ-bunches of Lie algebras.

Definition 2A. The *Lie* $\widetilde{\mathrm{m}}$YB-*algebra* is a Lie algebra $\mathfrak{g}$ with the bracket $[\cdot,\cdot]$, supplied by an operator $R:\mathfrak{g}\mapsto\mathfrak{g}$ such that

$$R[RX,Y]+R[X,RY]=[RX,RY]+R^2[X,Y].$$

Proposition 2. *The Jacobi identity holds for the bracket*

$$[X,Y]_R=[RX,Y]+[X,RY]-R[X,Y]$$

in any Lie $\widetilde{\mathrm{m}}$YB-*algebra.*

Theorem 1A states that any linear Γ-bunch of Lie algebras is a Lie $\widetilde{\mathrm{m}}$YB-algebra. The inverse statement is also true.

Theorem 1B. *Any Lie* $\widetilde{\mathrm{m}}$YB-*algebra* $(\mathfrak{g},R)$ *is supplied by a structure of a linear* Γ-*bunch of Lie algebras with brackets* $[\cdot,\cdot]_\lambda=[\cdot,\cdot]+\lambda[\cdot,\cdot]_R$ *and homomorphisms* $R_\lambda=1+\lambda R$. *In particular, the brackets* $[\cdot,\cdot]$ *and* $[\cdot,\cdot]_R$ *are compatible.*

Proposition 3. *For any Lie* $\widetilde{\mathrm{m}}$YB-*algebra* $(\mathfrak{g},R)$ *an operator* R' *of the form* $f(R)=a_0+a_1R+a_2R^2+\ldots+a_nR^n$ *defines a structure of the Lie* $\widetilde{\mathrm{m}}$YB-*algebra* $(\mathfrak{g},f(R))$.

Hence, the brackets $[\cdot,\cdot]_{R^n}$ $(n\in\mathbb{Z}_+)$ are compatible between each other. Proposition 3 allows to construct the operators R' such that $(R')^2=1$ for the subsequent applications of the factorization method.

1.4. Lie bi-$\widetilde{\mathrm{m}}$YB-algebras.

Proposition 4. *Let* $\mathfrak{A}$ *be an associative algebra then its commutator algebra* $\mathfrak{A}_{[\cdot,\cdot]}$ *is a Lie* $\widetilde{\mathrm{m}}$YB-*algebra, where* $R(X)=R^l_Q(X)=QX$ *or* $R(X)=R^r_Q(X)=XQ$ *are the operators of the left or right multiplication on the elements* Q *of the associative algebra* $\mathfrak{A}$. *Moreover,*

$$[X,Y]_{R^r_Q}=[X,Y]_{R^l_Q}=XQY-YQX.$$

This construction motivates the following definition.

Definition 2B. The *Lie bi*-$\widetilde{\mathrm{m}}$YB-*algebra* is a Lie algebra $\mathfrak{g}$ with bracket $[\cdot,\cdot]$ supplied by two commuting operators R_1 and R_2 such that $(\mathfrak{g},R_1)$ and $(\mathfrak{g},R_2)$ are the Lie $\widetilde{\mathrm{m}}$YB-algebras with identical brackets $[\cdot,\cdot]_{R_1}$ and $[\cdot,\cdot]_{R_2}$.

Thus proposition 4 states that the fixing of any element Q of the associative algebra $\mathfrak{A}$ supplies this algebra by a structure of the Lie bi-$\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{A}_{[\cdot,\cdot]},R^r_Q,R^l_Q)$.

Remark 4. In any Lie bi-$\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{g},R_1,R_2)$ the operator R_1–R_2 is a derivative of the bracket $[\cdot,\cdot]$ of the Lie algebra $\mathfrak{g}$.

Remark 5. A Lie $\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{g},R)$ is the Lie bi-$\widetilde{\mathrm{m}}$YB-algebra if and only if there exists a derivative ξ of the Lie algebra $\mathfrak{g}$ commuting with R such that

$$[\xi X,\xi Y]=[SX,Y]+[X,SY]-S[X,Y],\qquad S=R\xi,$$

and $R_1=R$, $R_2=R+\xi$.

Hence, in the Lie bi-$\widetilde{\mathrm{m}}$YB-algebra the operator ξ is a derivative of both brackets $[\cdot,\cdot]$ and $[\cdot,\cdot]_R$.

Proposition 5. *For any Lie bi-*$\widetilde{\mathrm{m}}$YB*-algebra* $(\mathfrak{g}, R_1, R_2)$ *and any polynomial* $f(x)$ *the triple* $(\mathfrak{g}, f(R_1), f(R_2))$ *is a Lie bi-*$\widetilde{\mathrm{m}}$YB*-algebra.*

Remark 6. In any Lie bi-$\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{A}_{[\cdot,\cdot]}, R^r_Q, R^l_Q)$ the bracket $[\cdot,\cdot]^q$ defined as

$$[X, Y]^q = [R^r_Q X, R^l_Q Y] + [R^l_Q X, R^r_Q Y] - R^r_Q R^l_Q [X, Y],$$

obeys the Jacobi identity

$$[[X, Y]^q, Z]^q + c.p. = 0,$$

and also is compatible with $[\cdot,\cdot]$, $[\cdot,\cdot]_{R_1}$ and $[\cdot,\cdot]_{R_2}$. Moreover, the identity

$$[X, Y]^q = [X, Y]_{R_1^2} = [X, Y]_{R_2^2}$$

holds for all X and Y.

This fact motivates the following definition.

Definition 2C. A Lie bi-$\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{g}, R_1, R_2)$ is called *even-tempered* if the identities

$$[R_1 X, R_2 Y] + [R_2 X, R_1 Y] - R_1 R_2 [X, Y] = [R_1^2 X, Y] + [X, R_1^2, Y] - R_1^2 [X, Y],$$

$$[R_1 X, R_2 Y] + [R_2 X, R_1 Y] - R_1 R_2 [X, Y] = [R_2^2 X, Y] + [X, R_2^2, Y] - R_2^2 [X, Y]$$

hold.

Note that one may claim only one of identities in definition 2C in view of proposition 5.

Remark 7. The identities in the even-tempered algebra have the form

$$[RX, \xi Y] + [\xi X, RY] - R\xi[X, Y] = [R^2 X, Y] - 2[RX, RY] + [X, R^2 Y]$$

in terms of R and ξ.

Thus, remark 6 means that the Lie bi-$\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{A}_{[\cdot,\cdot]}, R^r_Q, R^l_Q)$ constructed from an associative algebra $\mathfrak{A}$ and its element Q in proposition 4 is even-tempered.

2. Representations of linear bunches of Lie algebras

This paragraph is devoted to representations of linear bunches of Lie algebras, which naturally appear in the quantization of linearly depending on a parameter and compatible (linear) Poisson brackets (such situation is usual for systems in external magnetic fields [6]).

Definition 3 (cf.[3])**.** The *representation of the linear* $\Pi\Gamma$*-bunch* $\mathfrak{g}$ *of Lie algebras in the linear space* H is a mapping $T : \mathfrak{g} \mapsto \mathrm{End}(H)$ for which there exists an operator Q_R in H such that

$$T([X, Y]_\lambda) = T(X)(1 + \lambda Q_R)T(Y) - T(Y)(1 + \lambda Q_R)T(X)$$

for all X and Y from $\mathfrak{g}$. A linear $\Pi\Gamma$-bunch of Lie algebras is called *representable* if it admits a faithful representation.

The claim of a representability of a linear $\Pi\Gamma$-bunch is rather strong. Let us investigate some necessary conditions of representability of linear $\Pi\Gamma$-bunches.

Define for two Lie brackets $[\cdot,\cdot]_\alpha$ and $[\cdot,\cdot]_\beta$ in the linear space V their $\Diamond$-product (depending on the element of V) as

$$\begin{aligned}[X, Y]_{\alpha \underset{Z}{\Diamond} \beta} = & \frac{1}{2}\left([[X, Z]_\alpha, Y]_\beta + [[X, Y]_\alpha, Z]_\beta + [[Z, Y]_\alpha, X]_\beta - \right.\\ & \left. - [[X, Z]_\beta, Y]_\alpha - [[X, Y]_\beta, Z]_\alpha - [[Z, Y]_\beta, X]_\alpha\right).\end{aligned}$$

Let us formulate the main theorem on the necessary conditions of the representability of linear Γ-bunches of Lie algebras.

Theorem 2. *The linear* $\Pi\Gamma$*-bunch of Lie algebras* $\mathfrak{g}$ *is representable only if there exists a linear family of (compatible) Lie brackets, which contains the prescribed bunch and which is closed under the* $\Diamond$*-product. If such family* V *of Lie brackets exists then* $(\mathfrak{g}, V)$ *possesses a natural structure of an isotopic pair.*

One has the following corollary for linear $\Pi\Gamma$-bunches of Lie algebras.

Corollary. *A linear* Γ*-bunch of Lie algebras defined by the Lie* $\widetilde{\mathrm{m}}\mathrm{YB}$*-algebra* $(\mathfrak{g}, R)$ *is representable only if the pair* $(\mathfrak{g}, R + \lambda[\operatorname{ad} Z, R])$ *is a Lie* $\widetilde{\mathrm{m}}\mathrm{YB}$*-algebra for any element* Z *of the Lie algebra* $\mathfrak{g}$ *and an arbitrary number* λ*.*

Note that each homomorphism of the Lie $\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{g}, R)$ into the Lie bi-$\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{A}_{[\cdot,\cdot]}, R^r_Q, R^l_Q)$ of proposition 4 allows to construct representations of the linear Γ-bunch $\mathfrak{g}$ in the spaces of representations of the associative algebra $\mathfrak{g}$. The inverse statement is not true in general, i.e. a representation T of the linear Γ-bunch $\mathfrak{g}$ in the linear space H does not obligatory define a homomorphism of the Lie $\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{g}, R)$ into the Lie bi-$\widetilde{\mathrm{m}}$YB-algebra $(\mathrm{Mat}_{[\cdot,\cdot]}(n), R^r_Q, R^l_Q)$ (i.e. a homomorphism of $(\mathfrak{g}, R)$ into $(\mathrm{Mat}_{[\cdot,\cdot]}(n), R^l_Q)$ as algebras with operators) that is demonstarted by the following example.

Example 3. Let us consider the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ with the basis L_{-1}, L_0, L_1 and commutation relations $[L_i, L_j] = (i-j)L_{i+j}$ and the operator R on it: $RL_i = iL_i$. The pair $(\mathfrak{sl}(2,\mathbb{C}), R)$ is a Lie $\widetilde{\mathrm{m}}$YB-algebra and defines the linear Γ-bunch $\mathfrak{g}$. The two-dimensional fundamental representation of the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$ realizes a representation of this bunch with $Q = L_0$, however, none homomorphism of the Lie $\widetilde{\mathrm{m}}$YB-algebra $(\mathfrak{sl}(2,\mathbb{C}), R)$ into the Lie bi-$\widetilde{\mathrm{m}}$YB-algebra $(\mathrm{Mat}_{[\cdot,\cdot]}(n), R^r_Q, R^l_Q)$ corresponds to it, because the equality $T(R)T(L_0) = 0$ implies $T(R) = 0$ in view of the invertibility of the operator $T(L_0)$.

References

[1] *Semenov-Tyan-Shanskii M.A.*, What the classical r-matrix is. Funkts.anal.i ego prilozh. 17(4) (1983) 17-33.

[2] *Gelfand I.M., Dorfman I.Ya.*, Hamiltonian operators and classical Yang-Baxter equation. Funkts.anal.i ego prilozh. 16(4) (1982) 1-9.

[3] *Juriev D.V.*, Topics in isotopic pairs and their representations. I,II. Teor.Matem.Fiz. 105(1) (1995) 18-28; 111(1) (1997) 149-158.

[4] *Kon P.*, Universal algebra. Moscow, 1968.

[5] *Mal'tsev A.I.*, Algebraic systems. Moscow, 1970.

[6] *Dubrovin B.A., Krichever I.M., Novikov S.P.*, Integrable systems. In "Current Math. Problems. Basic Directions IV". Moscow, VINITI, 1988.

APPENDIX: THE ORIGINAL RUSSIAN VERSION OF ARTICLE

ИЗОТОПИЧЕСКИЕ ПАРЫ И ИХ ПРЕДСТАВЛЕНИЯ. III. ПУЧКИ АЛГЕБР ЛИ И МОДИФИЦИРОВАННОЕ УРАВНЕНИЕ ЯНГА-БАКСТЕРА

Д.В.Юрьев

Центр математическоё физики и информатики “Таласса Этерия”,
ул.Миклухо-Маклая 20-180, Москва 117437 Россия.
E-mail: denis@juriev.msk.ru

q-alg/9708027

Данная небольшая заметка является кратким комментарием к некоторым аспектам теории классическоё r-матрицы [1] и бигамильтонова формализма [2], мотивировкоё которого служат конструкции работ [3].

1. Линеёные пучки алгебр Ли и $\widetilde{\mathrm{m}}\mathrm{YB}$-алгебры Ли

Данныё параграф посвящен анализу некоторых алгебраических структур, связанных с линеёными пучками алгебр Ли.

1.1. Г-пучки алгебр Ли и касательная скобка.

Определение 1А. *Пучком (однопараметрическим) алгебр Ли* называется линеёное пространство $\mathfrak{g}$, снабженное однопараметрическим семеёством лиевских скобок $[\cdot,\cdot]_\lambda$ ($\lambda \in \mathbb{R}$). Алгебра Ли из пучка $\mathfrak{g}$, определяемая скобкоё $[\cdot,\cdot]_\lambda$, обозначается $\mathfrak{g}_\lambda$. Пучок алгебр Ли $\mathfrak{g}$ называется Γ-*пучком*, если заданы гомоморфизмы R_λ из алгебр Ли $\mathfrak{g}_\lambda$ в алгебру Ли $\mathfrak{g}_0$, иными словами, для всех X и Y из $\mathfrak{g}$ выполнено равенство

$$R_\lambda[X,Y]_\lambda = [R_\lambda X, R_\lambda Y],$$

где $[\cdot,\cdot] = [\cdot,\cdot]_0$.

В дальнеёшем будут рассматриваться гладкие пучки, т.е. пучки для которых лиевские скобки образуют гладкое семеёство, и гладкие Γ-пучки, для которых гладким является также семеёство гомоморфизмов R_λ.

Предложение 1. *Пусть $\mathfrak{g}$ - гладкиё Γ-пучок алгебр Ли, определим касательную скобку $[\cdot,\cdot]_R$ следующим образом:*

$$[X,Y]_R = \left.\frac{d[X,Y]_\lambda}{d\lambda}\right|_{\lambda=0}.$$

Тогда

$$[X,Y]_R = [RX,Y] + [X,RY] - R[X,Y],$$

где

$$R = \left.\frac{dR_\lambda}{d\lambda}\right|_{\lambda=0}.$$

Замехцание 1. Скобка $[\cdot,\cdot]_R$ тривиальна (тождественно равна нулю) тогда и только тогда, когда R – дифференцирование скобки $[\cdot,\cdot]$.

Замечание 2. Условие лиевости скобки $[\cdot,\cdot]_R$ имеет вид

$$\big([B(X,Y),Z]+B([X,Y],Z)\big)+c.p.=0,$$

где

$$B(X,Y)=R[RX,Y]+R[X,RY]-[RX,RY]-R^2[X,Y].$$

Развернутое выражение для $B(X,Y)$ может быть скомпоновано в двучленную формулу:

$$B(X,Y)=R[X,Y]_R-[RX,RY].$$

1.2. Линеёные Г-пучки алгебр Ли и модифицированное уравнение Янга-Бакстера. Отметим, что скобка $[\cdot,\cdot]$ и касательная скобка $[\cdot,\cdot]_R$ в произвольном Г-пучке алгебр Ли всегда удовлетворяет тождеству

$$\big([[X,Y]_R,Z]+[[X,Y],Z]_R\big)+c.p.=0.$$

Как следствие, если касательная скобка $[\cdot,\cdot]_R$ лиевская, то она согласована со скобкоё $[\cdot,\cdot]$. Это обстоятельство является мотивировкоё для рассмотрения линеёных Г-пучков алгебр Ли.

Пучок алгебр Ли $\mathfrak{g}$ называется линеёным, если лиевские скобки $[\cdot,\cdot]_\lambda$ образуют линеёное семеёство, иными словами $[\cdot,\cdot]_\lambda=(1-\lambda)[\cdot,\cdot]+\lambda[\cdot,\cdot]_1$. Г-пучок алгебр Ли называется линеёным, если не только лиевские скобки, но и гомоморфизмы R_λ образуют линеёное семеёство, т.е. $R_\lambda=1+\lambda R$. Требование линеёности семеёства гомоморфизмов R_λ естественно, поскольку для нелинеёного семеёства R_λ его линеёная часть $1+\lambda R$ ($R=\frac{dR_\lambda}{d\lambda}\big|_{\lambda=0}$) задает то же линеёное семеёство скобок $[\cdot,\cdot]_\lambda$.

Теорема 1А. *Для линеёного* Г*-пучка алгебр Ли* $\mathfrak{g}$ *касательная скобка* $[\cdot,\cdot]_R$ *(которая в этом случае лиевская) удовлетворяет условию*

$$R[X,Y]_R=[RX,RY],$$

иными словами,

$$R[RX,Y]+R[X,RY]=[RX,RY]+R^2[X,Y].$$

Теорема означает, что условие лиевости скобки $[\cdot,\cdot]_R$ из замечания 2 в случае линеёных Г-пучков факторизуется и сводится к требованию тождественного равенства выражения $B(X,Y)$ нулю. Отметим, что лиевость скобки $[\cdot,\cdot]_R$ в силу ее согласованности со скобкоё $[\cdot,\cdot]$ означает, что пучок $[\cdot,\cdot]+\lambda[\cdot,\cdot]_R$ – линеёныё пучок алгебр Ли, однако, он не обязан быть Г-пучком, поэтому в общем случае условие лиевости из замечания 2 не факторизуется.

Последнее тождество в теореме 1А совпадает с модифицированным классическим уравнением Янга-Бакстера [1] при $R^2=1$, что выполняется для наиболее важных примеров, рассмотренных в [1]. В этом случае, как показано в [1], лиевскоё скобкоё также является скобка $[\cdot,\cdot]'_R$, задаваемая следующим образом

$$[X,Y]'_R = [RX,Y] + [X,RY] = [X,Y]_R + R[X,Y].$$

Общее условие лиевости скобки $[\cdot,\cdot]'_R$ выглядит следующим образом:

$$[R^2[X,Y],Z] + c.p. = 0.$$

Скобка $[\cdot,\cdot]'_R$ играет ключевую рль в методе факторизации [1]. Приведем пример, когда $R^2 \neq 1$, однако, $[R^2[X,Y],Z] + c.p. = 0$.

Пример 1. Рассмотрим алгебру Витта с базисом e_k $(k \in \mathbb{Z})$ и коммутационными соотношениями

$$[e_i, e_j] = (i-j)e_{i+j}.$$

Введем операторы R_n следующего вида

$$R_n(e_i) = e_{i+n}.$$

Каждыё из операторов R_n задает на алгебре Витта структуру $\widetilde{\mathrm{m}}$YB-алгебры, при этом ни для какого n квадраты операторов R_n не равны единице, однако, при всех n тождество $[R_n^2[X,Y],Z] + c.p. = 0$ выполнено.

Замечание 3. Линеёные Г-пучки алгебр Ли не образуют подмногообразия (в смысле [4,5]) в многообразии всех линееных пучков алгебр Ли. Тождества, которые связывают скобки $[\cdot,\cdot]$ и $[\cdot,\cdot]_R$ (как, впрочем, и $[\cdot,\cdot]$ и $[\cdot,\cdot]'_R$) неизвестны.

Определение 1Б. Линеёныё пучок алгебр Ли $\mathfrak{g}$ называется *линеёным* ПГ-*пучком*, если он допускает мономорфизм в некоторыё линеёныё Г-пучок алгебр Ли.

Как отмечалось выше, описание класса линеёных ПГ-пучков алгебр Ли с помощью тождеств неизвестно.

Приведем пример линеёного ПГ-пучка алгебр Ли, не являющегося линеёным Г-пучком.

Пример 2 (ср.[3]). Пусть $\mathfrak{g} = \mathfrak{sa}(n)$ – алгебра Ли всех кососимметрических матриц $n \times n$ со стандартным коммутатором $[X,Y] = XY - YX$, и Q – произвольная нескалярная симметрическая матрица $n \times n$. Зададим семеёство согласованных лиевских скобок $[\cdot,\cdot]_\lambda$ следующим образом

$$[X,Y]_\lambda = [X,Y] + \lambda(XQY - YQX).$$

Полученныё линеёныё пучок алгебр Ли не является линеёным Г-пучком, поскольку в силу нескалярности матрицы Q некоторые из алгебр $\mathfrak{g}_\lambda$, будучи изоморфными $\mathfrak{sa}(p,q)$, не допускают гомоморфизмов в $\mathfrak{sa}(n)$. С другоё стороны, данныё пучок вкладывается в пучок всех матриц $n \times n$ с теми же скобками, которыё является линеёным Г-пучком с гомоморфизмами R_λ, задаваемыми операторами умножения справа или слева на матрицы $Q_\lambda = 1 + \lambda Q$.

1.3. $\widetilde{\mathrm{m}}\mathrm{YB}$-алгебры Ли и линейные Γ-пучки алгебр Ли.

Определение 2А. $\widetilde{\mathrm{m}}\mathrm{YB}$-*алгеброй Ли* называется алгебра Ли $\mathfrak{g}$ со скобкой $[\cdot,\cdot]$, снабженная оператором $R:\mathfrak{g}\mapsto\mathfrak{g}$ таким, что

$$R[RX,Y]+R[X,RY]=[RX,RY]+R^2[X,Y].$$

Предложение 2. *В* $\widetilde{\mathrm{m}}\mathrm{YB}$-*алгебре Ли для скобки*

$$[X,Y]_R=[RX,Y]+[X,RY]-R[X,Y]$$

выполняется тождество Якоби.

Теорема 1А утверждает, что всякий линейный Γ-пучок алгебр Ли является $\widetilde{\mathrm{m}}\mathrm{YB}$-алгеброй Ли. Имеет место и обратное утверждение.

Теорема 1Б. *Всякая* $\widetilde{\mathrm{m}}\mathrm{YB}$-*алгебра Ли* $(\mathfrak{g},R)$ *снабжается структурой линейного* Γ-*пучка алгебр Ли со скобками* $[\cdot,\cdot]_\lambda=[\cdot,\cdot]+\lambda[\cdot,\cdot]_R$ *и гомоморфизмами* $R_\lambda=1+\lambda R$. *В частности, скобки* $[\cdot,\cdot]$ *и* $[\cdot,\cdot]_R$ *согласованы.*

Предложение 3. *Для произвольной* $\widetilde{\mathrm{m}}\mathrm{YB}$-*алгебры Ли* $(\mathfrak{g},R)$ *любой оператор* R' *вида* $f(R)=a_0+a_1R+a_2R^2+\ldots+a_nR^n$ *задает структуру* $\widetilde{\mathrm{m}}\mathrm{YB}$-*алгебры Ли* $(\mathfrak{g},f(R))$.

Как следствие, скобки $[\cdot,\cdot]_{R^n}$ $(n\in\mathbb{Z}_+)$ согласованы между собой. Предложение 3 позволяет строить операторы R' такие, что $(R')^2=1$, для последующего применения метода факторизации.

1.4. Би-$\widetilde{\mathrm{m}}\mathrm{YB}$-алгебры Ли.

Предложение 4. *Пусть* $\mathfrak{A}$ *– ассоциативная алгебра, тогда коммутаторная алгебра* $\mathfrak{A}_{[\cdot,\cdot]}$ *является* $\widetilde{\mathrm{m}}\mathrm{YB}$-*алгеброй Ли, где* $R(X)=R^l_Q(X)=QX$ *или* $R(X)=R^r_Q(X)=XQ$ *– операторы умножения слева и справа на элементы* Q *ассоциативной алгебры* $\mathfrak{A}$. *При этом,*

$$[X,Y]_{R^r_Q}=[X,Y]_{R^l_Q}=XQY-YQX.$$

Эта конструкция является мотивировкой для следующего определения.

Определение 2Б. *Би-*$\widetilde{\mathrm{m}}\mathrm{YB}$-*алгеброй Ли* называется алгебра Ли $\mathfrak{g}$ со скобкой $[\cdot,\cdot]$, снабженная двумя коммутирующими операторами R_1 и R_2 такими, что $(\mathfrak{g},R_1)$ и $(\mathfrak{g},R_2)$ – $\widetilde{\mathrm{m}}\mathrm{YB}$-алгебры Ли с совпадающими скобками $[\cdot,\cdot]_{R_1}$ и $[\cdot,\cdot]_{R_2}$.

Таким образом, предложение 4 утверждает, что фиксирование произвольного элемента Q ассоциативной алгебры $\mathfrak{A}$ снабжает указанную алгебру структурой би-$\widetilde{\mathrm{m}}\mathrm{YB}$-алгебры Ли $(\mathfrak{A}_{[\cdot,\cdot]},R^r_Q,R^l_Q)$.

Замечание 4. Оператор R_1-R_2 в би-$\widetilde{\mathrm{m}}\mathrm{YB}$-алгебре $(\mathfrak{g},R_1,R_2)$ является дифференцированием скобки $[\cdot,\cdot]$ алгебры Ли $\mathfrak{g}$.

Замечание 5. $\widetilde{\mathrm{m}}\mathrm{YB}$-алгебра Ли $(\mathfrak{g},R)$ является би-$\widetilde{\mathrm{m}}\mathrm{YB}$-алгеброй тогда и только тогда, когда существует дифференцирование ξ алгебры Ли $\mathfrak{g}$, коммутирующее с R, такое, что

$$[\xi X,\xi Y]=[SX,Y]+[X,SY]-S[X,Y],\qquad S=R\xi.$$

При этом $R_1=R$, $R_2=R+\xi$.

Как следствие, оператор ξ в би-$\widetilde{\mathrm{m}}\mathrm{YB}$-алгебре Ли является дифференцированием обеих скобок $[\cdot,\cdot]$ и $[\cdot,\cdot]_R$.

Предложение 5. *Для любоё би-*$\widetilde{\mathrm{m}}$YB*-алгебры Ли* $(\mathfrak{g}, R_1, R_2)$ *и многочлена* $f(x)$ *троёка* $(\mathfrak{g}, f(R_1), f(R_2))$ *является би-*$\widetilde{\mathrm{m}}$YB*-алгеброё Ли.*

Замечание 6. В би-$\widetilde{\mathrm{m}}$YB-алгебре Ли $(\mathfrak{A}_{[\cdot,\cdot]}, R^r_Q, R^l_Q)$ скобка $[\cdot,\cdot]^q$, задаваемая следующим образом

$$[X,Y]^q = [R^r_Q X, R^l_Q Y] + [R^l_Q X, R^r_Q Y] - R^r_Q R^l_Q [X,Y],$$

удовлетворяет тождеству Якоби

$$[[X,Y]^q, Z]^q + c.p. = 0,$$

а также согласована со скобками $[\cdot,\cdot]$, $[\cdot,\cdot]_{R_1}$ и $[\cdot,\cdot]_{R_2}$. Более того, для всех X и Y выполнено равенство

$$[X,Y]^q = [X,Y]_{R_1^2} = [X,Y]_{R_2^2}.$$

Этот факт является мотивировкоё для следующего определения.

Опредление 2B. Би-$\widetilde{\mathrm{m}}$YB-алгебра Ли $(\mathfrak{g}, R_1, R_2)$ называется *уравновешенноё*, если выполнены следующие тождества

$$\begin{aligned}
&[R_1X, R_2Y] + [R_2X, R_1Y] - R_1R_2[X,Y] = [R_1^2X, Y] + [X, R_1^2, Y] - R_1^2[X,Y],\\
&[R_1X, R_2Y] + [R_2X, R_1Y] - R_1R_2[X,Y] = [R_2^2X, Y] + [X, R_2^2, Y] - R_2^2[X,Y].
\end{aligned}$$

Отметим, что в определении 2B достаточно потребовать выполнения одного из тождеств в силу предложения 5.

Замечание 7. Тождества уравновешенности в терминах R и ξ имеют вид

$$[RX, \xi Y] + [\xi X, RY] - R\xi[X,Y] = [R^2X, Y] - 2[RX, RY] + [X, R^2Y].$$

Таким образом, замечание 6 означает, что би-$\widetilde{\mathrm{m}}$YB-алгебра $(\mathfrak{A}_{[\cdot,\cdot]}, R^r_Q, R^l_Q)$, построенная в предложении 4 по ассоциативноё алгебре $\mathfrak{A}$ и ее элементу Q, является уравновешенноё.

2. Представления линеёных пучков алгебр Ли

Данныё параграф посвящен представлениям линеёных пучков алгебр Ли, которые естественно возникают в рамках квантования линеёно зависящих от параметра согласованных (линеёных) скобок Пуассона (что характерно для систем во внешнем магнитном поле [6]).

Определение 3 (ср.[3]). *Представлением линеёного* ПГ*-пучка* $\mathfrak{g}$ *алгебр Ли в линеёном пространстве* H, называется отображение $T : \mathfrak{g} \mapsto \mathrm{End}(H)$, для которого существует оператор Q_R в H такоё, что

$$T([X,Y]_\lambda) = T(X)(1+\lambda Q_R)T(Y) - T(Y)(1+\lambda Q_R)T(X),$$

для произвольных X и Y из $\mathfrak{g}$. Линеёныё ПГ-пучок алгебр Ли называется *представимым*, если он обладает хотя бы одним точным представлением.

Требование представимости линейного ПГ-пучка достаточно сильное. Изучим ряд необходимых условий представимости линейных ПГ-пучков.

Определим для двух лиевских скобок $[\cdot,\cdot]_\alpha$ и $[\cdot,\cdot]_\beta$, заданных в линейном пространстве V, их $\Diamond$-произведение (зависящее от элемента пространства V) следующим образом:

$$[X,Y]_{\alpha\underset{Z}{\Diamond}\beta}=\frac{1}{2}\left([[X,Z]_\alpha,Y]_\beta+[[X,Y]_\alpha,Z]_\beta+[[Z,Y]_\alpha,X]_\beta-\right.$$
$$\left.-[[X,Z]_\beta,Y]_\alpha-[[X,Y]_\beta,Z]_\alpha-[[Z,Y]_\beta,X]_\alpha\right).$$

Сформулируем основную теорему о необходимых условиях представимости линейных Г-пучков алгебр Ли.

Теорема 2. *Линейный* ПГ*-пучок алгебр Ли* $\mathfrak{g}$ *представим, только если существует линейное семейство (совместимых) лиевских скобок, содержащее указанный пучок и замкнутое относительно* $\Diamond$*-произведения. Если такое семейство* V *лиевских скобок существует, то* $(\mathfrak{g},V)$ *наделяется естественной структурой изотопической пары.*

Для линейных ПГ-пучков алгебр Ли имеем следующее следствие.

Следствие. *Линейный* Г*-пучок алгебр Ли, задаваемый* $\widetilde{\mathrm{m}}$YB*-алгеброй Ли* $(\mathfrak{g},R)$*, представим, только если для любого элемента* Z *алгебры Ли* $\mathfrak{g}$ *и произвольного числа* λ *пара* $(\mathfrak{g},R+\lambda[\mathrm{ad}\,Z,R])$ *является* $\widetilde{\mathrm{m}}$YB*-алгеброй Ли.*

Отметим, что по всякому гомоморфизму $\widetilde{\mathrm{m}}$YB-алгебры Ли $(\mathfrak{g},R)$ в би-$\widetilde{\mathrm{m}}$YB-алгебру Ли $(\mathfrak{A}_{[\cdot,\cdot]},R_Q^r,R_Q^l)$ предложения 4 строятся представления линейного Г-пучка $\mathfrak{g}$ в пространствах представлений ассоциативной алгебры $\mathfrak{g}$. Обратное, вообще говоря, не верно, т.е. представление T линейного Г-пучка $\mathfrak{g}$ в линейном пространстве H не всегда определяет гомоморфизм $\widetilde{\mathrm{m}}$YB-алгебры Ли $(\mathfrak{g},R)$ в би-$\widetilde{\mathrm{m}}$YB-алгебру Ли $(\mathrm{Mat}_{[\cdot,\cdot]}(n),R_Q^r,R_Q^l)$ (т.е. гомоморфизм $(\mathfrak{g},R)$ в $(\mathrm{Mat}_{[\cdot,\cdot]}(n),R_Q^l)$ как алгебр с операторами), что показывает следующий пример.

Пример 3. Рассмотрим алгебру Ли $\mathfrak{sl}(2,\mathbb{C})$ с базисом L_{-1}, L_0, L_1 и коммутационными соотношениями $[L_i,L_j]=(i-j)L_{i+j}$, а также оператор R на неё: $RL_i=iL_i$. Пара $(\mathfrak{sl}(2,\mathbb{C}),R)$ является $\widetilde{\mathrm{m}}$YB-алгеброй Ли и определяет линейный Г-пучок $\mathfrak{g}$. Фундаментальное двумерное представление алгебры Ли $\mathfrak{sl}(2,\mathbb{C})$ осуществляет представление этого пучка с $Q=L_0$, однако, ему не соотвецтвует никакой гомоморфизм $\widetilde{\mathrm{m}}$YB-алгебры Ли $(\mathfrak{sl}(2,\mathbb{C}),R)$ в би-$\widetilde{\mathrm{m}}$YB-алгебру Ли $(\mathrm{Mat}_{[\cdot,\cdot]}(n),R_Q^r,R_Q^l)$ поскольку в силу обратимости оператора $T(L_0)$ равенство $T(R)T(L_0)=0$ влечет $T(R)=0$.

Список литературы

[1] *Семенов-Тян-Шанский М.А.*, Что такое классическая r-матрица // Функц.анал. и его прилож. 1983. Т.17(4). С.17-33.

[2] *Гельфанд И.М., Дорфман И.Я.*, Гамильтоновы операторы и классическое уравнение Янга-Бакстера // Функц.анал.и его прилож. 1982. Т.16(4). С.1-9.

[3] *Юрьев Д.В.*, Изотопические пары и их представления. I,II // ТМФ. 1995. Т.105(1). С.18-28; ТМФ. 1997. Т.111(1). С.149-158.

[4] *Кон П.*, Универсальная алгебра. М., 1968.

[5] *Мальцев А.И.*, Алгебраические системы. М., 1970.

[6] *Дубровин Б.А., Кричевер И.М., Новиков С.П.*, Интегрируемые системы / Современные пробл.математики. Фундам. направления IV. М., ВИНИТИ, 1988.